

A Local-Global Criterion for Dynamics on $\mathbb{P}^1$

JOSEPH H. SILVERMAN AND JOSÉ FELIPE VOLOCH

ABSTRACT. Let K be a number field or a one-dimensional function field, let $\varphi : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ be a rational map of degree at least two defined over K , let $P \in \mathbb{P}^1(K)$ be a point with infinite φ -orbit, and let $Z \subset \mathbb{P}^1$ be a finite set of points. We prove a local-global criterion for the intersection of the φ -orbit of P and the finite set Z . This is a special case of a dynamical Brauer-Manin criterion suggested by Hsia and Silverman.

INTRODUCTION

A recent paper of Hsia and Silverman [3] describes an analog of the Brauer–Manin obstruction in the setting of arithmetic dynamics. The definition is based on work of Scharaschkin [6], who showed how to reformulate the Brauer–Manin obstruction on curves of genus at least 2 as a purely adèlic-geometric statement, with no reference to cohomology. (See also [5].) In this note we prove a dynamical analog of Scharaschkin’s conjecture for dynamical systems on $\mathbb{P}^1$. The proof uses a variety of tools, including a Zsigmondy theorem for primitive divisors in dynamical systems that was recently proven by Ingram and Silverman [4].

We recall from [3] the setup for the dynamical version of the Brauer–Manin obstruction.

Notation. Let K be a number field, let X/K be a projective variety, and let

$$\varphi : X \rightarrow X$$

be a K -morphism of infinite order. Let $\mathbf{A}_K$ denote the ring of adèles of K , and for any point $P \in X(K)$, write $\mathbf{C}(\mathcal{O}_\varphi(P))$ for the closure

Date: June 16, 2008.

1991 *Mathematics Subject Classification.* Primary: 11B37; Secondary: 11G99, 14G99, 37F10.

Key words and phrases. arithmetic dynamical systems, local-global criterion, Brauer–Manin obstruction.

The first author’s research supported by NSF DMS-0650017.

of the orbit $\mathcal{O}_\varphi(P)$ of P in $X(\mathbf{A}_K)$. Let Z be a subvariety of X . We clearly have an inclusion

$$\mathcal{O}_\varphi(P) \cap Z(K) \subset \mathcal{C}(\mathcal{O}_\varphi(P)) \cap Z(\mathbf{A}_K). \quad (1)$$

The primary question raised in [3] is to ask when the inclusion (1) is an equality.

More generally, we can ask a similar question after excluding a (finite) set of places. For any set of places S we let

$$\mathbf{A}'_{K,S} = \prod'_{v \notin S} K_v \subset \mathbf{A}_K,$$

where the product is the usual adelic restricted product, and we write $\mathcal{C}_S(\mathcal{O}_\varphi(P))$ for the closure of the orbit in $\mathbf{A}'_{K,S}$.

Our main result is a proof of a local-global equality (1) for $X = \mathbb{P}^1$.

Theorem 1. *Let K be a number field, let S be a finite set of places of K , and let $\varphi : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ be a rational map of degree at least two defined over K . Let $Z \subset \mathbb{P}^1(K)$ be a finite set of points. Then with notation as above,*

$$\mathcal{O}_\varphi(P) \cap Z(K) = \mathcal{C}_S(\mathcal{O}_\varphi(P)) \cap Z(\mathbf{A}'_{K,S}).$$

Remark 2. As noted in [3], if (1) is to be an equality, one should require that the subvariety Z contains no positive dimensional φ -preperiodic subvarieties. However, since our subvariety Z is a set of points, this condition on Z is vacuous.

In Section 2 we prove an analogous result for function fields K/k . If the map φ is not isotrivial, the proof is essentially the same as the proof of Theorem 1, but the isotrivial case must be handled separately. Finally, in Section 3, we make some further comments and raise some related questions.

1. LOCAL-GLOBAL DYNAMICS ON $\mathbb{P}^1$ OVER NUMBER FIELDS

In this section we prove Theorem 1. We briefly recall some basic definitions from dynamical systems.

Definition. Let $\varphi(z) \in K(z)$ be a rational function of degree $d \geq 2$, which we may view as a morphism $\varphi : \mathbb{P}^1_K \rightarrow \mathbb{P}^1_K$. A point $\gamma \in \mathbb{P}^1(\bar{K})$ is *periodic* for φ if $\varphi^n(\gamma) = \gamma$ for some $n \geq 1$. The smallest such n is called the φ -*period* of γ . We say that γ is *preperiodic* for φ if its φ -*orbit*

$$\mathcal{O}_\varphi(\gamma) = \{\gamma, \varphi(\gamma), \varphi^2(\gamma), \dots\}$$

is finite. Equivalently, γ is preperiodic if some iterate $\varphi^n(\gamma)$ is periodic.

Remark 3. During the proof of Theorem 1, we make use of the following notation. For any two points $x, y \in \mathbb{P}^1(K)$ and any prime ideal $\mathfrak{p}$ of K , we write

$$x \equiv y \pmod{\mathfrak{p}} \quad (2)$$

to indicate that the reductions of x and y modulo $\mathfrak{p}$ coincide. Formally, this notation means that if we write $x = [x_1, x_2]$ and $y = [y_1, y_2]$ using $\mathfrak{p}$ -integral coordinates such that at least one coordinate is a $\mathfrak{p}$ -unit, then

$$x_1y_2 - x_2y_1 \equiv 0 \pmod{\mathfrak{p}}.$$

Alternatively, we define a (nonarchimedean) chordal metric on $\mathbb{P}^1$ for the absolute value v by (cf. [11, §2.1])

$$\Delta_v(x, y) = \frac{|x_1y_2 - x_2y_1|_v}{\max\{|x_1|_v, |x_2|_v\} \cdot \max\{|y_1|_v, |y_2|_v\}}.$$

This is independent of the choice of homogeneous coordinates for x and y , and we define the congruence (2) to mean $\Delta_v(x, y) < 1$, where v is the absolute value associated to the prime $\mathfrak{p}$. More generally, we write

$$\prod (\alpha_i - \beta_i) \equiv 0 \pmod{\mathfrak{p}}$$

for points $\alpha_i, \beta_i \in \mathbb{P}^1(K)$ to mean that $\prod \Delta_v(\alpha_i, \beta_i) < 1$.

Proof of Theorem 1. Without loss of generality, we may assume that the set S contains all archimedean places of K and all nonarchimedean places at which φ has bad reduction (see [11, §2.5]). Let $\beta \in \mathbb{P}^1(K)$. If β is preperiodic, then its orbit $\mathcal{O}_\varphi(\beta)$ is finite, so the orbit is discrete in $\mathbb{P}^1(K_v)$ for every place v . Hence in this case we have $\mathcal{C}(\mathcal{O}_\varphi(\beta)) = \mathcal{O}_\varphi(\beta)$, so the equality

$$\mathcal{O}_\varphi(\beta) \cap Z(K) = \mathcal{C}(\mathcal{O}_\varphi(\beta)) \cap Z(\mathbf{A}_K).$$

is obvious.

We suppose now that $\beta \in \mathbb{P}^1(K)$ is a point with infinite φ -orbit. Since Z is a finite set, there can be only finitely many iterates $\varphi^n(\beta)$ that lie in Z , so replacing β with some $\varphi^n(\beta)$, we may assume that $\mathcal{O}_\varphi(\beta) \cap Z(K) = \emptyset$. Under this assumption, we suppose that there exists a point

$$\alpha \in \mathcal{C}_S(\mathcal{O}_\varphi(\beta)) \cap Z(\mathbf{A}'_{K,S})$$

and derive a contradiction. The supposed existence of α means that there is an increasing sequence of positive integers

$$\mathcal{N} = \{n_1, n_2, n_3, \dots\}$$

such that the sequence of points $\varphi^n(\beta)$ with $n \in \mathcal{N}$ converges adelically to $\alpha \in Z(\mathbf{A}'_{K,S})$. Looking at each coordinate of the adele α , we see in

particular that for every prime $\mathfrak{p} \notin S$ of K there exists an integer $N_{\mathfrak{p}}$ such that

$$\varphi^n(\beta) \equiv \alpha_{\mathfrak{p}} \pmod{\mathfrak{p}} \quad \text{for all } n \in \mathcal{N} \text{ with } n \geq N_{\mathfrak{p}}.$$

Note that since Z is a finite subset of $\mathbb{P}^1(K)$, each $\alpha_{\mathfrak{p}}$ is a point in $\mathbb{P}^1(K)$ considered as a subset of $\mathbb{P}^1(K_{\mathfrak{p}})$.

We are going to use the dynamical Zsigmondy theorem from [4], whose statement we recall after making one definition.

Definition. Let $\varphi : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ and let $\gamma \in \mathbb{P}^1$. We say that φ is of *polynomial type at γ* if there is some $k \geq 1$ such that γ is a totally ramified fixed point of φ^k .

Remark 4. It is easy to check that φ is of polynomial type at γ if and only if, when we conjugate φ by $(z - \gamma)^{-1}$, the map φ^k is conjugated to a polynomial in z . We also remark that it is an exercise using the Riemann-Hurwitz genus formula to show that if φ is of polynomial type at γ , then the φ -period of γ is at most 2, cf. [11, Theorem 1.7].

Theorem 5. (Ingram–Silverman [4]) *Let K be a number field, let S be a finite set of primes, let $\beta \in \mathbb{P}^1(K)$ be a point with infinite orbit, and let $\gamma \in \mathbb{P}^1(K)$ be a preperiodic point. Let $\varphi : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ be a K -morphism of degree at least two that is not of polynomial type at γ . There exists an integer $M = M_{\varphi, \beta, \gamma}$ so that for all $m \geq M$ there exists a prime ideal $\mathfrak{q}_m \notin S$ of K satisfying*

$$\begin{aligned} \varphi^m(\beta) &\equiv \gamma \pmod{\mathfrak{q}_m}, \\ \varphi^i(\beta) &\not\equiv \gamma \pmod{\mathfrak{q}_m} \quad \text{for all } 0 \leq i < m. \end{aligned}$$

In the terminology of [4], the prime ideal $\mathfrak{p}_m$ is a *primitive divisor* for $\varphi^m(\beta) - \gamma$. Theorem 5 says that $\varphi^m(\beta) - \gamma$ has a primitive divisor for all sufficiently large values of m . There are many classical theorems showing the existence of primitive divisors for the multiplicative group and for elliptic curves, see for example [7, 8, 12]. The proof of Theorem 5 uses a dynamical analog [10] of Siegel's theorem [9, IX.3.1] for integral points on elliptic curves.

Remark 6. The theorem stated in [4] does not exclude a finite set of primes, but since S is finite, we immediately get Theorem 5 from [4] by increasing M so as to exclude the finitely many m such that the primitive divisor of $\varphi^m(\beta) - \gamma$ is in S .

We fix an integer $R \geq 3$ and extend the field K so that all of the periodic points of φ of exact period R are in $\mathbb{P}^1(K)$. Let $\gamma \in \mathbb{P}^1(K)$ be a point of exact period R for φ . As noted in Remark 4, the map φ

is not of polynomial type at γ , so we can apply Theorem 5 to find an integer $M = M_{\varphi, \beta, \gamma}$ such that $\varphi^m(\beta) - \gamma$ has a non- S primitive divisor for all $m \geq M$. Doing this for each of the finitely many points of period R , we may choose one $M = M_{\varphi, \beta, R}$ that works for all γ of period R .

Next we fix an integer $T \geq 1$ and we choose a non- S primitive divisor for $\varphi^m(\beta) - \gamma$ for each m between M and $M+T$. Thus we find *distinct* prime ideals $\mathfrak{q}_{M, \gamma}, \mathfrak{q}_{M+1, \gamma}, \dots, \mathfrak{q}_{M+T, \gamma} \notin S$ such that

$$\varphi^m(\beta) \equiv \gamma \pmod{\mathfrak{q}_{m, \gamma}} \quad \text{for } M \leq m \leq M+T.$$

We have indicated the dependence of $\mathfrak{q}_{m, \gamma}$ on both m and γ , because we now replace γ by $\varphi^i \gamma$ for each $1 \leq i < R$ and apply the same argument. Thus we find prime ideals $\mathfrak{q}_{m, \varphi^i \gamma} \notin S$ satisfying

$$\varphi^m(\beta) \equiv \varphi^i(\gamma) \pmod{\mathfrak{q}_{m, \varphi^i \gamma}} \quad \text{for } M \leq m \leq M+T \text{ and } 0 \leq i < R.$$

Further, for any particular value of i , the ideals $\mathfrak{q}_{m, \varphi^i \gamma}$ are distinct for $M \leq m \leq M+T$. (These ideals also depend on φ and β , of course, but φ and β are fixed throughout the proof.) This gives us a finite set of non- S primes

$$\mathcal{Q} = \mathcal{Q}_{R, T} = \{\mathfrak{q}_{m, \varphi^i \gamma} : M \leq m \leq M+T \text{ and } 0 \leq i < R\}.$$

(Note that $M = M_{\varphi, \beta, R}$ is independent of T .)

Consider the product

$$\prod_{r=0}^{R-1} (\varphi^m(\beta) - \varphi^r(\gamma)). \quad (3)$$

It is divisible by each of the (not necessarily distinct) prime ideals

$$\mathfrak{q}_{m, \gamma}, \mathfrak{q}_{m, \varphi \gamma}, \mathfrak{q}_{m, \varphi^2 \gamma}, \dots, \mathfrak{q}_{m, \varphi^{R-1} \gamma}. \quad (4)$$

Further, the fact that γ is periodic with period R implies that for any $n \geq m$, the product

$$\prod_{r=0}^{R-1} (\varphi^n(\beta) - \varphi^r(\gamma)). \quad (5)$$

is divisible by all of the primes in the list (4). To see this, let $\mathfrak{q}$ be any prime listed in (4). Then we use the R -periodicity of γ to write

$$\prod_{r=0}^{R-1} (\varphi^n(\beta) - \varphi^r(\gamma)) = \prod_{r=0}^{R-1} (\varphi^n(\beta) - \varphi^{r+1}(\gamma)),$$

and this last product is clearly divisible by

$$\prod_{r=0}^{R-1} (\varphi^{n-1}(\beta) - \varphi^r(\gamma)).$$

(We are using the general fact that $\varphi(b) - \varphi(a)$ is divisible by $b - a$.) Now a downward induction shows that (5) is divisible by (3).

Recall the sequence of positive integer $\mathcal{N}$ such that $\varphi^n(\beta)$ converges $\mathbf{A}'_{K,S}$ -adically to α as $n \rightarrow \infty$ with $n \in \mathcal{N}$. In particular, for each $\mathfrak{q} \in \mathcal{Q}$ we have

$$\varphi^n(\beta) \equiv \alpha_{\mathfrak{q}} \pmod{\mathfrak{q}} \quad (6)$$

for all sufficiently large $n \in \mathcal{N}$. (Note that $\mathcal{Q} \cap S = \emptyset$.) We recall that each $\alpha_{\mathfrak{q}}$ is chosen from the finite set Z , so we can reformulate (6) as

$$\prod_{a \in Z} (\varphi^n(\beta) - a) \equiv 0 \pmod{\mathfrak{q}} \quad (7)$$

for sufficiently large $n \in \mathcal{N}$ and for every $\mathfrak{q} \in \mathcal{Q}$.

For notational convenience, we define

$$\mathfrak{Q}_{R,T} = \prod_{\mathfrak{q} \in \mathcal{Q}_{R,T}} \mathfrak{q} = \text{Radical} \left(\prod_{M \leq m \leq M+T} \prod_{0 \leq i < R} \mathfrak{q}_{m, \varphi^i \gamma} \right).$$

The fact that the ideals $\mathfrak{q}_{m, \varphi^i \gamma}$ are distinct for fixed i and varying m implies that

$$\#\mathcal{Q}_{R,T} > T,$$

so $\mathfrak{Q}_{R,T}$ is a product of more than T distinct prime ideals.

With this notation, we can rewrite (7) as

$$\prod_{a \in Z} (\varphi^n(\beta) - a) \equiv 0 \pmod{\mathfrak{Q}_{R,T}} \quad \text{for sufficiently large } n \in \mathcal{N}. \quad (8)$$

Similarly, it follows from our earlier discussion that

$$\prod_{r=0}^{R-1} (\varphi^n(\beta) - \varphi^r(\gamma)) \equiv 0 \pmod{\mathfrak{Q}_{R,T}} \quad \text{for all } n \geq M_R. \quad (9)$$

Comparing (8) and (9) for any single sufficiently large value of $n \in \mathcal{N}$, we conclude that

$$\prod_{r=0}^{R-1} \prod_{a \in Z} (\varphi^r(\gamma) - a) \equiv 0 \pmod{\mathfrak{Q}_{R,T}}. \quad (10)$$

This last congruence is very interesting, because the product on the left-hand side does not depend on T , while the ideal $\mathfrak{Q}_{R,T}$ is a product of more than T distinct prime ideals. Letting $T \rightarrow \infty$, we conclude that

the product in (10) vanishes. Therefore there is at least one value r_0 and at least one point $a_0 \in Z$ such that

$$\varphi^{r_0}(\gamma) = a_0.$$

To recapitulate, we have proven that given any point $\gamma \in \mathbb{P}^1$ that is periodic for φ of period at least 3, there is some point in the φ -orbit of γ that lies in the set Z . But a rational map on $\mathbb{P}^1$ has periodic points of infinitely many periods (more precisely, it has a periodic point of exact period R for every $R \geq 5$, see [2]), which contradicts the assumption that Z is a finite set. $\square$

2. LOCAL-GLOBAL DYNAMICS ON $\mathbb{P}^1$ OVER FUNCTION FIELDS

In this section we prove the analogue of Theorem 1 for function fields. Let k be an algebraically closed field and let K/k be a function field, that is, K is a finitely generated extension of k of transcendence degree one. We employ the same notation as in the introduction except that we consider only the places of K/k , i.e., the absolute values of K that are trivial on k .

Theorem 7. *Let K/k be a function field as above, let S be a finite set of places of K/k , and let $\varphi : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ be a rational map of degree at least two defined over K . Let $Z \subset \mathbb{P}^1(K)$ be a finite set of points. Then with notation as in the introduction,*

$$\mathcal{O}_\varphi(P) \cap Z(K) = \mathcal{C}_S(\mathcal{O}_\varphi(P)) \cap Z(\mathbf{A}'_{K,S}).$$

Proof. A rational map φ defined over K is isotrivial (or split) if there exists a Möbius transformation $M \in \mathrm{PGL}_2(\bar{K})$ such that $M \circ \varphi \circ M^{-1}$ has coefficients in k . As pointed out by T. Tucker [4, Remark 4], the Zsigmondy theorem in [4] can be proved for non-isotrivial rational maps over function fields using the results of Baker [1]. We observe that the characteristic zero assumption is not used in [1], and therefore is not needed to extend the results of [4], contrary to what is stated there. Thus the argument used to prove Theorem 1 applies verbatim in the function field case if φ is non-isotrivial.

Assume now that φ is isotrivial. There is no loss of generality in extending K by adjoining the coefficients of M , and then replacing φ by $M \circ \varphi \circ M^{-1}$, we may assume that φ has coefficients in k .

Arguing again as in the proof of Theorem 1, the case of preperiodic points is straightforward.

We suppose now that $\beta \in \mathbb{P}^1(K)$ is a point with infinite φ -orbit. Since Z is a finite set, there can be only finitely many iterates $\varphi^n(\beta)$ that lie in Z , so replacing β with some $\varphi^n(\beta)$, we may assume that

$\mathcal{O}_\varphi(\beta) \cap Z = \emptyset$. Under this assumption, we suppose that there exists a point

$$\alpha \in \mathbb{C}_S(\mathcal{O}_\varphi(\beta)) \cap Z(\mathbf{A}'_{K,S})$$

and derive a contradiction. The supposed existence of α means that there is an increasing sequence of positive integers

$$\mathcal{N} = \{n_1, n_2, n_3, \dots\}$$

such that the sequence of points $\varphi^n(\beta)$ with $n \in \mathcal{N}$ converges adelically to $\alpha \in Z(\mathbf{A}'_{K,S})$.

We note first that $\beta \notin k$, since otherwise the sequence $\varphi^{n_i}(\beta) \in k$ could not converge $\mathfrak{p}$ -adically for any place $\mathfrak{p}$ of K/k unless it became eventually constant, which would force β to be preperiodic, contrary to our current assumption.

Assume now that φ is not purely inseparable. Choose an element $b \in k$ that is periodic with respect to φ and such that the backwards orbit of b with respect to φ is infinite and such that no element of Z is in the (forward) orbit of b . It follows that the set T of places $\mathfrak{p}$ of K/k for which an element of Z is congruent modulo $\mathfrak{p}$ to an element of the (forward) orbit of b is finite. Now choose an element $a \in k$ in the backwards orbit of b with respect to φ , such that a is not in the (forward) orbit of b and such that the places of K/k extending the place $\beta - a$ of $k(\beta)$ do not belong to T . Let $\mathfrak{p}$ be a place of K/k extending the place $\beta - a$ of $k(\beta)$. By assumption, there exists $\alpha_{\mathfrak{p}} \in Z(K)$ with $\varphi^{n_i}(\beta)$ converging $\mathfrak{p}$ -adically to $\alpha_{\mathfrak{p}}$. Thus, for all large i , we have

$$\varphi^{n_i}(a) \equiv \varphi^{n_i}(\beta) \equiv \alpha_{\mathfrak{p}} \pmod{\mathfrak{p}}.$$

But, for i large, $\varphi^{n_i}(a)$ belongs to the forward orbit of b . This forces $\mathfrak{p} \in T$, which is a contradiction.

Finally, we have to deal with the case that φ is purely inseparable. Without loss of generality, we may assume that $\varphi(z) = z^q$, where q is a power of the characteristic of K . Let $a \in k$ and let $\mathfrak{p}$ be a place of K/k extending the place $\beta - a$ of $k(\beta)$. Then the only way that a sequence $\varphi^{n_i}(\beta)$ can converge $\mathfrak{p}$ -adically is if a is in a finite field, and in this case converges to an element in the Frobenius orbit of a . There are only finitely many such a whose orbits intersect the finite set Z , and hence $\varphi^{n_i}(\beta)$ cannot converge adelically. This completes the proof. $\square$

3. COMMENTS AND SPECULATIONS

Remark 8. We have stated our principal results (Theorems 1 and 7) for rational maps of degree at least two, but we can also consider maps of degree one. After a finite extension of the field K , any map $\varphi : \mathbb{P}^1 \rightarrow \mathbb{P}^1$

of degree one may be conjugated to a map either of the form $\varphi(z) = az$ or of the form $\varphi(z) = z + 1$. There are three cases to consider.

- (1) If $\varphi(z) = az$ and a is a root of unity, say $a^n = 1$, then $\varphi^n(z) = z$ is the identity map, so every point has finite orbit. Hence the closure of $\mathcal{O}_\varphi(P)$ is equal to $\mathcal{O}_\varphi(P)$, so the conclusions of Theorems 1 and 7 are clearly true.
- (2) If $\varphi(z) = az$ and a is not a root of unity, then we can mimic the proof of Theorem 1, replacing Theorem 5 with the classical Zsigmondy theorem for the multiplicative group. So Theorems 1 and 7 are also true in this case.
- (3) If $\varphi(z) = z + 1$, then Theorem 1 is not true in general. For example, take $Z = \{0, \infty\}$ and $\beta = 1$. Then the orbit of β is $\mathcal{O}_\varphi(\beta) = \{1, 2, 3, \dots\}$, which does not intersect Z . On the other hand, the orbit of β contains the subsequence $\{n! : n = 1, 2, 3, \dots\}$, and this subsequence converges to the point in $Z(\mathbf{A}_\mathbb{Q})$ that is ∞ at the archimedean place and is 0 at all nonarchimedean places. Hence $\mathcal{C}(\mathcal{O}_\varphi(\beta)) \cap Z(\mathbf{A}_\mathbb{Q}) \neq \emptyset$.

Remark 9. Let $K/\mathbb{Q}$ be a number field, let $f(z) \in K(z)$ be a rational function of degree at least two, and let $\varphi(z) = z - f(z)/f'(z)$. Iteration of $\varphi(z)$ is Newton's method of finding a root of $f(z)$. Let $\alpha \in K$ be a point whose orbit $\mathcal{O}_\varphi(\alpha)$ does not contain a root of f , e.g., any point that is not preperiodic for φ . Then Theorem 1 says that there are infinitely many places $v \in S$ such that Newton's method applied to α does not converge to a root of $f(z)$.

We raise the following question: In this situation (with $\deg f(z) \geq 3$), is it true that Newton iteration applied to α converges for infinitely many places and also does not converge for infinitely many places?

Remark 10. We observe that an equality

$$\mathcal{O}_\varphi(P) \cap Z(K) = \mathcal{C}(\mathcal{O}_\varphi(P)) \cap Z(\mathbf{A}_K),$$

such as given in [3] and in Theorem 1, provides an algorithm for determining if $\mathcal{O}_\varphi(P) \cap Z(K)$ is nonempty. Thus “by day” one computes elements of the orbit and checks if they are in Z , while “by night” one checks if there are points in $\mathcal{O}_\varphi(P) \cap Z$ modulo $\mathfrak{Q}$ for ideals $\mathfrak{Q}$ that are more and more divisible by primes not in S . For a fixed $\mathfrak{Q}$ this is a finite computation. This is analogous to the fact that the Scharaschkin conjecture [6] gives an algorithm to decide whether a curve of genus at least 2 has any rational points (but not necessarily to find them all).

Acknowledgements. The authors would like to thank the organizers of the TateFest (May 2008), at which event this research was initiated.

REFERENCES

- [1] M. Baker. A finiteness theorem for canonical heights attached to rational maps over function fields, *J. Reine Angew. Math.*, to appear. ArXiv math.NT/0601046v2.
- [2] A. F. Beardon. *Iteration of Rational Functions*, volume 132 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1991.
- [3] L.-C. Hsia and J. H. Silverman. On a dynamical Brauer–Manin obstruction. Proceedings of JA2007, to appear.
- [4] P. Ingram and J. H. Silverman. Primitive divisors in arithmetic dynamics. *Proc. Camb. Philos. Soc.*, 2008. to appear.
- [5] B. Poonen and J. F. Voloch. The Brauer–Manin obstruction for subvarieties of abelian varieties over function fields. ArXiv math.NT/0612263, December 2006.
- [6] V. Scharaschkin. *Local-global problems and the Brauer–Manin obstruction*. PhD thesis, University of Michigan, 1999.
- [7] A. Schinzel. Primitive divisors of the expression $A^n - B^n$ in algebraic number fields. *J. Reine Angew. Math.*, 268/269:27–33, 1974.
- [8] J. H. Silverman. Wieferich’s criterion and the *abc*-conjecture. *J. Number Theory*, 30(2):226–237, 1988.
- [9] J. H. Silverman. *The Arithmetic of Elliptic Curves*, volume 106 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1992.
- [10] J. H. Silverman. Integer points, Diophantine approximation, and iteration of rational maps. *Duke Math. J.*, 71(3):793–829, 1993.
- [11] J. H. Silverman. *The Arithmetic of Dynamical Systems*, volume 241 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2007.
- [12] K. Zsigmondy. Zur Theorie der Potenzreste. *Monatsh. Math. Phys.*, 3(1):265–284, 1892.

MATHEMATICS DEPARTMENT, BOX 1917, 151 THAYER STREET, BROWN UNIVERSITY, PROVIDENCE, RI 02912 USA

E-mail address: jhs@math.brown.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TEXAS, 1 UNIVERSITY STATION C1200, AUSTIN, TX 78712 USA

E-mail address: voloch@math.utexas.edu